

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2012 (19.04.2012)

PCT

(10) International Publication Number
WO 2012/051116 A1

- (51) **International Patent Classification:**
A01N 37/30 (2006.01) *A61K 31/205* (2006.01)
- (21) **International Application Number:**
PCT/US2011/055617
- (22) **International Filing Date:**
10 October 2011 (10.10.2011)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/392,294 12 October 2010 (12.10.2010) US
- (71) **Applicant (for all designated States except US):** THE MEDICINES COMPANY [US/US]; 8 Slyvan Way, Parsippany, NJ 07054 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** RAJESHWAR, Motheram [US/US]; 17 Summerfield Blvd., Dayton, NJ 08810 (US). WILLIAMS, Gregory, Charles [US/US]; 50 Old Colony Road, Bernardsville, NJ 07924 (US).
- (74) **Agents:** WANG, Lisa et al.; Gibbons P.C, One Gateway Center, Newark, NJ 07102 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))



WO 2012/051116 A1

(54) **Title:** CLEVIDIPINE EMULSION FORMULATIONS CONTAINING ANTIMICROBIAL AGENTS

(57) **Abstract:** Pharmaceutical formulations comprising clevidipine and an antimicrobial agent exhibit a reduced propensity for microbial growth and provide increased convenience to health care workers administering clevidipine-containing formulations to patients.

CLEVIDIPINE EMULSION FORMULATIONS CONTAINING ANTIMICROBIAL AGENTS

Field of the Invention

[0001] The instant invention relates to a stable, pharmaceutical oil-in-water emulsion formulation for parenteral administration that includes clevidipine and an antimicrobial agent.

Background of the Invention

[0002] Clevidipine is a dihydropyridine calcium channel blocker that reduces blood pressure in a subject to which it is administered. It is characterized as a short-acting, highly selective drug that is typically used in a hospital setting due to its rapid metabolism, exhibiting an initial phase half-life of about one minute and a terminal half-life of approximately 15 minutes. More detailed information on short-acting dihydropyridines can be found in U.S. Patent No. 5,856,346, the entire disclosure of which is incorporated by reference herein as if set forth in its entirety.

[0003] Clevidipine is further characterized by having low solubility in water and moderate to high solubility in lipids. When clevidipine is dissolved in oil-in-water emulsions, it results in better solubility and/or less side effects than when other conventional solution formulations are utilized. Cleviprex® is a clevidipine oil-in-water emulsion formulation approved by the U.S. FDA for intravenous administration in the treatment of acute hypertension, primarily in the emergency room and intensive care unit, and in pre-, peri-, and post-operative settings.

[0004] Cleviprex® can support microbial growth since it contains soybean oil and egg yolk phospholipids. Therefore, clevidipine oil-in-water emulsion formulations require strict aseptic technique during handling and administration to avoid microbial contamination that can cause infections among patients. To minimize the possibility of microbial contamination, it is recommended that such formulations be discarded four hours after opening. Such a requirement places a burden on health care providers in that fresh vials of the drug must be continually obtained and set up while the patient is being treated. Accordingly, there exists a need for a stable clevidipine emulsion formulation that possesses greater antimicrobial properties, thereby diminishing the risk of microbial contamination in patients and providing greater ease in handling. Such a formulation would also result in cost savings to the health care providers and patients in decreasing the wastage of clevidipine and reducing the time-consuming efforts involved with manipulation and replacement of vials containing the drug.

Summary of the Invention

[0005] Based on the state of the art, it is the object of the present invention to provide a stable clevidipine emulsion formulation with a reduced propensity for microbial contamination.

[0006] In a first embodiment of the present invention, there is provided a pharmaceutical formulation comprising clevidipine, or a pharmaceutically acceptable salt or ester thereof, an antimicrobial agent, a lipid, an emulsifier, a tonicity modifier and water.

[0007] In one aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water. The tonicity of emulsion is adjusted with glycerin and further comprises an amount of EDTA sufficient to inhibit growth of microorganisms in the formulation in the event of accidental extrinsic contamination.

[0008] In second aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water. The tonicity of emulsion is adjusted with glycerin and further comprises an amount of EDTA and sodium citrate sufficient to inhibit growth of microorganisms in the formulation in the event of accidental extrinsic contamination.

[0009] In a second embodiment of the invention, there is provided a pharmaceutical formulation comprising clevidipine, or a pharmaceutically acceptable salt or ester thereof, an antimicrobial agent, a lipid, an emulsifier, a co-emulsifier, a tonicity modifier and water.

[0010] In one aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water, and wherein the emulsion is further stabilized by the co-emulsifier oleic acid. The tonicity of emulsion is adjusted with glycerin and further comprises an amount of EDTA sufficient to inhibit growth of microorganisms in the formulation.

[0011] In a second aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water, and wherein the emulsion is further stabilized by the co-emulsifier oleic acid. The tonicity of emulsion is adjusted with glycerin and

further comprises an amount of EDTA and sodium citrate sufficient to inhibit growth of microorganisms in the formulation in the event of accidental extrinsic contamination.

[0012] In a third embodiment of the invention, there is provided a pharmaceutical formulation comprising clevidipine, or a pharmaceutically acceptable salt or ester thereof, an antimicrobial agent, an antioxidant, a lipid, an emulsifier, a tonicity modifier and water.

[0013] In one aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water and stabilized by the antioxidant sodium ascorbate. The tonicity of emulsion is adjusted with glycerin and further comprises an amount of EDTA sufficient to inhibit growth of microorganisms in the formulation.

[0014] In a fourth embodiment of the invention, there is provided a pharmaceutical formulation comprising clevidipine, or a pharmaceutically acceptable salt or ester thereof, an antimicrobial agent, an antioxidant, a lipid, an emulsifier, a co-emulsifier, a tonicity modifier and water.

[0015] In one aspect of this embodiment, the present invention provides an oil-in-water emulsion wherein clevidipine is dispersed or dissolved in a lipid, such as soy bean oil, which in turn is emulsified with egg yolk phospholipids in water and wherein the emulsion is further physically stabilized by the co-emulsifier oleic acid and chemically stabilized by the antioxidant sodium ascorbate. The tonicity of emulsion is adjusted with glycerin and further comprises an amount of EDTA sufficient to inhibit growth of microorganisms in the formulation.

[0016] In aspects of embodiments where the formulation comprises an antioxidant, the antioxidant may reduce the formation of oxidative degradants H324/78 and H152/66

[0017] In preferred aspects of each embodiment of the invention, the antimicrobial agent is an alcohol or a chelating agent. More preferably, the antimicrobial agent is disodium edetate (EDTA) or sodium citrate, or both.

[0018] In preferred aspects of each embodiment of the invention, the pharmaceutical formulation is sterile. In the event of accidental contamination, the pharmaceutical formulation will retard the growth of microorganisms to no more than 1 log within at least about 24 hours.

[0019] In each embodiment of the invention, clevidipine and the emulsion maintain their stability in the formulation. In each embodiment of the invention, the pharmaceutical formulation is for parenteral administration. In each embodiment of the invention, the pharmaceutical formulation will have a pH that ranges from about 6.0 to about 8.8.

Brief Description of the Figures

[0020] Understanding of the present invention will be facilitated by consideration of the following detailed description of the embodiments of the present invention taken in conjunction with the accompanying drawings:

[0021] Figure 1: clevidipine degradation pathway.

Detailed Description of the Invention

[0022] It is to be understood that the figures and descriptions of the present invention have been simplified to illustrate elements that are relevant for a clear understanding of the present invention, while eliminating, for the purpose of clarity, many other elements found in typical pharmaceutical compositions and methods of stabilization. Those of ordinary skill in the art will recognize that other elements and/or steps are desirable and/or required in implementing the present invention. However, because such elements and steps are well known in the art, and because they do not facilitate a better understanding of the present invention, a discussion of such elements and steps is not provided herein. The disclosure herein is directed to all such variations and modifications to such elements and methods known to those skilled in the art. Furthermore, the embodiments identified and illustrated herein are for exemplary purposes only, and are not meant to be exclusive or limited in their description of the present invention.

[0023] As used herein, the term “**clevidipine**” shall mean and include all varieties or forms of clevidipine. Unless otherwise specified, examples of such forms include all pharmaceutically acceptable salts, esters, isomers, stereo isomers, crystalline and amorphous forms. One particular example is clevidipine butyrate. The amount of clevidipine in the formulations of the present invention can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of clevidipine in the formulation

will generally range from about 0.005 to about 1.0% w/v, and includes ranges of from about 0.03 to about 0.5% w/v, and from about 0.01 to about 1.0% w/v. In particular embodiments, the amount of clevidipine in the formulation will be about 0.05, 0.1, or 0.3% w/v.

[0024] As used herein, the term “**pharmaceutically acceptable salt**” shall refer to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Examples of salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganese salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

[0025] As used herein, the term “**pharmaceutically acceptable ester**” shall refer to esters prepared by reaction of an oxo-acid with a compound containing a hydroxyl group. Generally esters are derived from an inorganic or organic acid and an alcohol. More commonly esters are prepared by condensing an organic acid and an alcohol. Examples of suitable esters that may be used in the formulations of the present invention include butyric acid esters, such as those prepared in accordance with teachings in U.S. Patent Nos. 5856346, 5739152, 6350877 and the like.

[0026] Clevidipine is manufactured by reaction of 4-(2',3'-dichlorophenyl)-1,4-dihydro-5-methoxycarbonyl-2,6-dimethyl-3-pyridinecarboxylic acid with chloromethyl butyrate to obtain clevidipine. This reaction can be done optionally in the presence of a corresponding hydrogen carbonate, such as KHCO_3 , in refluxing acetonitrile. Inorganic salts can be removed by filtration and the product is crystallized by the addition of isopropanol and water with subsequent cooling. It can also be crystallized by exchanging solvent from acetonitrile to a mixture of alcohol, such

as ethanol or isopropanol, and water with repeated evaporations. In the further purification of the product the crystals are washed with a mixture of water and ethanol or isopropanol. The product can be dissolved in refluxing isopropanol, crystallized by cooling, isolated by filtration and finally washed with a water and isopropanol mixture. A more detailed description of the manufacturing process of clevidipine can be found in U.S. Patent No. 6,350,877, the entire disclosure of which is incorporated by reference herein as if set forth in its entirety. As used herein, the range of clevidipine that may be used in the formulations includes a range of from about 0.005% to about 1% w/v.

[0027] Compositions containing clevidipine as an active ingredient are sensitive to water, heat, and light. Clevidipine degrades under adverse conditions into several impurities that compromise the potency of clevidipine. A degradation pathway of clevidipine is shown in Figure 1. This pathway identifies a number of clevidipine degradation products, including for example, H152/81, H168/79, H207/59, H324/78 and H152/66.

[0028] As with any of the exemplary embodiments of pharmaceutical compositions described herein, it is preferred that the level of clevidipine impurities in the composition is as low as possible. Therefore, while various exemplary embodiments of pharmaceutical compositions include levels of impurities within acceptable and effective ranges for the compositions as a whole, the more pure the composition, the more desirable the composition. The amount of clevidipine degradation products present in the formulations of the present invention is less than about 5%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, or 0.2%, or even less.

[0029] The pharmaceutical formulations of the present invention are emulsion formulations. For compounds that have poor solubility and stability in water, emulsions offer much better solubility, and stability than conventional aqueous solutions. Oil-in-water emulsions also prevent the compound from adherence to plastic infusion sets that are to be used when administering the compound. Further information regarding the formulation of clevidipine can be found in U.S. Patent No. 5,739,152, the entire disclosure of which is incorporated by reference herein as if set forth in its entirety.

[0030] As used herein, the term “**antimicrobial agent**” means an agent that inhibits the growth of microorganisms such as bacteria and fungi (molds and yeast). Particular classes of antimicrobial agents include chelating agents and alcohols. Chelating agents include, but are not

limited to, ethylenediamine tetraacetic acid (EDTA) and salts thereof, citric acid and salts thereof, and the like. Exemplary chelating agents are disodium edetate and sodium citrate. Alcohols include, but are not limited to, benzyl alcohol and chlorobutanol. Representative antimicrobial agents include EDTA, ascorbic acid, BHA/BHT, benzyl alcohol, benzoic acid, citric acid, edetic acid, parabens, phenol, propyl gallate, sorbic acid, sodium bisulfite, sodium sulfite, benzoic acid, chlorobutanol, chlorocresol, cresol, dehydroacetic acid, phenol, potassium benzoate, potassium sorbate, sodium benzoate, sodium dehydroacetate, sodium propionate, sorbic acid, thymol, benzalkonium chloride, benzethonium chloride, butyl paraben, cetylpyridinium chloride, ethylparaben, methylparaben, methylparaben sodium, propylparaben, propylparaben sodium, chlorocresol, cresol, dehydroacetic acid, ethylparaben, methylparaben, methylparaben sodium, phenol, potassium sorbate, thimersol and the various salt forms for these compounds.

[0031] In general the term “EDTA” means ethylenediamine tetraacetic acid and its derivatives and salts thereof. Particular derivative of use in the present invention includes disodium edetate. The nature of the EDTA derivative or salt is not critical, provided that it inhibits growth of microorganisms in the formulations of the present invention.

[0032] The amount of antimicrobial agent in the formulations of the present invention can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of the antimicrobial agent in the formulation will generally range from about 0.001 to about 1.5% w/v, and includes from about 0.005 to about 0.5% w/v, from about 0.001 to about 0.1% w/v, from about 0.01 to about 0.1% w/v, and from about 0.1 to about 1% w/v. In particular embodiments, the amount of the antimicrobial agent in the formulation will be about 0.1, 0.2, or 0.4% w/v. Where a chelating agent is used as the antimicrobial agent, the amount of chelating agent in the formulation will generally range from about 0.001% to about 0.5% w/v. The antimicrobial agent may comprise one, or more than one agent, such as two, three, four, etc., different antimicrobial agents.

[0033] As used herein, the term “antioxidant” in the formulations includes sodium ascorbate, cysteine hydrochloride, sodium bisulfite, sodium metabisulfite, sodium sulfite, ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), propyl gallate, tocopherol, and their pharmaceutically acceptable salts. An exemplary antioxidant is sodium ascorbate. The amount of antioxidant in the formulations of the present invention can vary

depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of the antioxidant in the formulation will generally range from about 0.01 to about 1.0% w/v, and includes from about 0.05 to about 1.0% w/v, and from about 0.05 to about 0.5% w/v. In particular embodiments, the amount of antioxidant in the formulation will be about 0.1% w/v.

[0034] As used herein, the term “**lipid**” in the formulations is any pharmaceutically acceptable oil, preferably a triglyceride such as soybean oil, safflower seed oil, olive oil, cottonseed oil, sunflower oil, sesame oil, peanut oil, corn oil, a medium chain triglyceride (such as Miglyol™ 812 or 810) or triacetin. The lipid may also be a propylene glycol diester or monoglyceride (such as acetylareal monoglyceride). The lipid can also be a mixture of one or more of these lipids. An exemplary lipid is soybean oil. The amount of lipid in the formulations of the present invention can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of lipid in the formulation will generally range from about 2 to about 30% w/v, and includes from about 5 to about 30% w/v, and from about 10% to about 20% w/v. In particular embodiments, the amount of lipid in the formulation will be about 20% w/v.

[0035] As used herein, the term “**emulsifier**” represents suitable pharmaceutically acceptable surfactants used in the formulations, preferably naturally occurring phospholipids extracted from egg yolk or soy bean, synthetic phosphatidyl cholines or purified phosphatidyl cholines from vegetable origin. Hydrogenated derivatives can also be used, such as phosphatidyl choline hydrogenated (egg) and phosphatidyl choline hydrogenated (soya). The amount of phospholipid emulsifier in the formulations of the present invention can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of emulsifier in the formulation will generally range from about 0.2 to about 2.0% w/v, and includes from about 0.5 to about 1.5% w/v. In particular embodiments, the amount of emulsifier in the formulation will be about 1.2 % w/v.

[0036] As used herein, the term “**co-emulsifier**” represents a second pharmaceutically acceptable surfactant that may be included in the formulations of the invention. Such surfactants include synthetic nonionic surfactants such as poloxamers (for example Poloxamer 188 and 407), Cremophor™, poloxamines, polyoxyethylene stearates, polyoxyethylene sorbitan fatty acid

esters or sorbitan fatty acid esters, derivatives of tocopherol such as tocopherol PEG succinate, long chain fatty acids such as oleic acid, stearic acid, palmitic acid, bile acids such as cholic acid and deoxycholic acid or surface active derivatives, and pharmaceutically acceptable salts thereof. An exemplary surfactant is oleic acid. The amount of co-emulsifier in the formulations of the present invention, when present, can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of surfactant in the formulation will generally range from about 0.005 to about 2% w/v, and includes from about 0.01 to about 2% w/v, and from about 0.01 to about 1.0% w/v. In particular embodiments, the amount of surfactant in the formulation will be about 0.03% w/v.

[0037] As used herein, the term “**tonicity modifier**” in the formulations of the present invention includes sodium chloride, potassium chloride, mannitol, sucrose, lactose, fructose, maltose, dextrose, dextrose anhydrous, propylene glycol, glycerol and glycerin. An exemplary tonicity modifier is glycerin. The amount of tonicity modifier in the formulations of the present invention can vary depending on the total overall volume of the formulation and the concentration of the other components. However, the amount of tonicity modifier in the formulation will generally range from about 2 to about 3% w/v, and includes from about 2 to about 2.75% w/v. In particular embodiments, the amount of tonicity modifier in the formulation will be about 2.25% w/v. The terms “tonicity modifier” and “isotonicity adjuster” are used herein interchangeably.

[0038] The amount of water in the formulations of the present invention, such as water-for-injections, is used to make up the volume to 100% w/v and can vary depending on the total overall volume of the formulation and the concentration of the other components.

[0039] As the formulations are intended for parenteral administration, the skilled artisan will understand that one or more additional components used in parenteral formulations may be included. Such additional components include stabilizing agents (e.g. carbohydrates, amino acids and polysorbates, such as 5% dextrose), solubilizing agents (e.g. cetrimide, sodium docusate, glyceryl monooleate, polyvinylpyrrolidone (PVP) and polyethylene glycol (PEG), buffers (e.g. acetates, citrates, phosphates, tartrates, lactates, succinates, amino acids and the like), preservatives (e.g. BHA, BHT, gentisic acids, vitamin E, ascorbic acid, sodium ascorbate and sulfur containing agents such as sulfites, bisulfites, metabisulfites, thioglycerols,

thioglycolates and the like), suspending or viscosity agents, chelating agents, and administration aids (e.g. local anesthetics, anti-inflammatory agents, anti-clotting agents, vaso-constrictors for prolongation and agents that increase tissue permeability).

[0040] Parenteral modes of administration include without limitation, intradermal, subcutaneous (s.c., s.q., sub-Q, Hypo), intramuscular (i.m.), intravenous (i.v.), intraperitoneal (i.p.), intra-arterial, intramedullary, intracardiac, intra-articular (joint), intrasynovial (joint fluid area), intracranial, intraspinal, and intrathecal (spinal fluids). Any known device useful for parenteral injection or infusion of drug formulations can be used to effect such administration.

[0041] In intravenous use, a sterile formulation of the present invention can be dissolved or suspended in any of the commonly used sterile intravenous fluids and administered by infusion. Intravenous fluids include, without limitation, physiological saline, phosphate buffered saline, 5% dextrose in water or Ringer's™ solution. The parenteral dosage form of formulations of the present invention can also be a ready-to-use solution in sterile sealed vials, hermetically sealed ampoules or in sterile pre-filled syringes, for example.

[0042] The sterile pre-filled syringes are syringes that contain a unit dose of a pharmaceutical formulation of the present invention. Suitable syringes are widely available and well known to the skilled artisan. An exemplary sterile pre-filled syringe is one that has been loaded with a unit dose of the pharmaceutical formulation and that is enclosed in an opaque, sealed package from which oxygen has been excluded. For example, oxygen may be displaced with CO₂ and/or N₂.

[0043] The pharmaceutical formulations of the present invention will have a pH that ranges from about 6.0 to about 8.8. In particular embodiments, the pH ranges from about 6.5 to about 8.0. In particular embodiments, the pH is 6.2, 6.5, 6.75, 7.0, or 7.5. Bases such as NaOH, KOH, and Ca(OH)₂ may be used to achieve the desired pH.

[0044] As used herein, the terms "inhibit", "inhibiting" and "inhibition" have their ordinary and customary meanings, and include inhibiting the growth of a bacteria or fungus in the formulations of the present invention. Such inhibition may be described as no more than about 10 fold growth for at least 24 hours following a low level (1 – 1000 Cfu/mL) of extrinsic contamination. Such growth may be determined, for example, by determining the number of colony forming units in the formulation when cultured at room temperature.

[0045] The duration of time over which inhibition of microbial growth is maintained will vary depending on the environmental conditions to which the formulation is exposed, e.g., the conditions under which a sterile vial of the formulation is pierced by a needle or sterility is otherwise breached. However, in one embodiment of the invention, microbial growth is inhibited for at least about 24 or more hours after the formulation is exposed to low level extrinsic microbial contamination.

[0046] The skilled artisan will understand that the pharmaceutical formulations of the present invention may be prepared using art-accepted means for preparing emulsion formulations. A general procedure for preparing clevidipine formulations is described as follows: an oil phase containing soybean oil, clevidipine and egg yolk phospholipids is mixed with an aqueous phase containing glycerin, at approximately 70° C to form a coarse emulsion. The pH of the coarse emulsion is adjusted using sodium hydroxide. Following pH adjustment, the coarse emulsion is homogenized under high pressure to produce a fine particle size and thus a stable emulsion. The emulsion is filled into appropriate containers and sterilized in an autoclave.

[0047] The invention is described in greater detail by reference to the following non-limiting examples.

Examples 1-5

[0048] **Example 1** - Formulations containing antimicrobial agents as shown in Table 1 below were prepared by adding stock solutions of anti microbial agent at appropriate concentration to clevidipine emulsions.

Table 1

		(% w/v)				
Function		Example 1	Example 2	Example 3	Example 4	Example 5
Active ingredient	Clevidipine butyrate	0.05	0.05	0.05	0.05	0.05
Oil phase	Soybean oil	20	20	20	20	20
Isotonicity adjuster	Glycerin	2.25	2.25	2.25	2.25	2.25
Microbial Growth Retardant	Disodium Edetate	0	0.005	0.01	0	0.005
Microbial Growth Retardant	Benzyl Alcohol	0	0	0	0.4	0.1
Emulsifier	Purified egg yolk phospholipids	1.2	1.2	1.2	1.2	1.2
pH adjuster	Sodium hydroxide	q.s	q.s	q.s	q.s	q.s
Aqueous Phase	Water for Injection	q.s to 100%	q.s to 100%	q.s to 100%	q.s to 100%	q.s to 100%

[0049] Sterile clevidipine oil-in-water compositions prepared as described above were evaluated for microbial growth. Microbial growth inhibition was tested using standard UPS test organisms. Each formulation was tested against five standard USP test organisms: *S. aureus* (SA, ATCC # 6538), *P. aeruginosa* (PA, ATCC #9027), *E. coli* (EC, ATCC # 8739), *C. albicans* (CA, ATCC # 10231), and *A. niger* (AN, ATCC #16404). Formulations were inoculated with suspensions of each organism in an amount that recovers less than 100 CFU (colony forming units) per filter. Inoculations for each formulation were conducted in triplicate.

[0050] Test samples were stored at 20-25° C for the duration of the study. Testing was performed at Time 0, 24, and 48 hours after inoculation. At appropriate time points, the samples were filtered through a 0.45 micrometer membrane filter and washed with 0.1% sterile peptone solution to remove the preservatives for the purpose of recovering the microorganisms. The washed filters were aseptically removed and placed on appropriate microbiological growth medium.

[0051] *S. aureus*, *P. aeruginosa* and *E. coli* were plated on tripticase soy agar and incubated aerobically at 30-35° C for 2 days or until countable colonies were formed, whichever occurred first. *C. albicans* and *A. niger* were plated on sabouraud dextrose agar and aerobically incubated at 20-25° C for 7 days or until countable colonies were formed, whichever occurred first.

[0052] Microbial inhibition was considered acceptable if there was a delay in onset or retardation of growth such that there was less than 10-fold (1 log) increase in viable colonies over a 24-hour period.

[0053] The effectiveness of EDTA alone or in combination with benzyl alcohol on in microbial inhibition over a 24 hour period is demonstrated in Table 2 below. From these experiments it can be concluded that the presence of EDTA and benzyl alcohol either alone or in combination demonstrated microbial retardation capabilities.

Table 2

Formulation	Microbial Retardant (% w/v)	Time Point	EC	SA	PA	CA	AN
Example 1	None	0	21	32	7	29	14
		24 hours	TNTC	> 300	210	TNTC	19
		48 hours	C	TNTC	C	C	35
Example 2	0.005% EDTA	0	30	24	11	35	36
		24 hours	2	21	2	233	30
		48 hours	0.3	14	5	310	25
Example 3	0.01% EDTA	0	23	5	5	20	31
		24 hours	4	15	0.1	77	17
		48 hours	3	16	0	250	40
Example 4	0.4% Benzyl Alcohol	0	26	21	17	22	16
		24 hours	63	34	2	45	19
		48 hours	102	66	1	910	37
Example 5	0.1% Benzyl Alcohol /0.005% EDTA	0	26	19	15	34	38
		24 hours	0.3	10	0.3	300	36
		48 hours	2	5	0	370	51

TNTC = too numerous to count

C = Confluent growth

[0054] Example 2 - Formulations containing antimicrobial agents as shown in Table 3 below were prepared by adding stock solutions of anti microbial agent at appropriate concentration to clevipidine emulsions.

Table 3

		(%w/v)	
Function		Example 6	Example 7
Active ingredient	Clevidipine butyrate	0.05	0.05
Oil phase	Soybean oil	20	20
Isotonicity adjuster	Glycerin	2.25	2.25
Microbial Growth Retardant	Disodium Edetate	0.005	0.005
Microbial Growth Retardant	Sodium Citrate	0.1	0
Antioxidant	Sodium Ascorbate	0	0.1
Emulsifier	Purified egg yolk phospholipids	1.2	1.2
pH adjuster	Sodium hydroxide	q.s	q.s
Aqueous Phase	Water for Injection	q.s to 100%	q.s.to 100%

[0055] Compositions described above were evaluated for microbial growth following the procedure described previously. The effectiveness of EDTA in combination with sodium citrate or sodium ascorbate in retarding microbial growth over a 24 hour period is demonstrated in Table 4 below. From these experiments it can be concluded that the presence of EDTA in combination with sodium citrate or sodium ascorbate demonstrated microbial retardation capabilities.

Table 4

Formulation	Microbial Retardant (% w/v)	Time Point	EC	SA	PA	CA	AN
Example 1	None	0	21	32	7	29	14
		24 hours	TNTC	> 300	210	TNTC	19
		48 hours	C	TNTC	C	C	35
Example 6	0.005% EDTA / 0.1% Sodium Citrate	0	19	14	9	25	31
		24 hours	2	5	4	55	39
		48 hours	0	3	23	85	34
Example 7	0.005% EDTA / 0.1% Sodium Ascorbate	0	25	30	16	29	34
		24 hours	0	25	26	209	37
		48 hours	0	19	42	3 >300	45

TNTC = too numerous to count

C = Confluent growth

[0056] **Example 3** - Clevidipine emulsion formulations were exposed to high intensity light (combination of cool white fluorescent and near UV light) in an effort to demonstrate the impact of light on the chemical stability and the results are shown in Table 5 below:

Table 5

Formulation	Microbial Retardant (% w/v)	Antioxidant (%w/v)	Oxidative Degradants Light Exposure (1.2 Million Lux Hours)	
			H324/78	H152/66
Example 1	None	None	0.2	0.4
Example 2	0.005% EDTA	None	0.2	0.2
Example 7	0.005% EDTA	0.1% Sodium Ascorbate	< 0.1	Not Detected

It was unexpectedly found that the addition of ascorbic acid in the presence of EDTA was effective in inhibition of photodegradation of clevidipine in the emulsion, without counteracting the antimicrobial effectiveness of EDTA.

[0057] **Example 4** - Clevidipine emulsion formulations with EDTA alone or in combination with sodium citrate were prepared by adding EDTA and/or sodium citrate as a part of the emulsion process in order to confirm the antimicrobial characteristics previously observed when added as a solution to a prepared emulsion. The emulsion formulations shown below were prepared as follows: The oil phase containing soybean oil and egg yolk phospholipids was mixed with the aqueous phase containing glycerin, disodium edetate with or without sodium citrate and water at approximately 70° C to form a coarse emulsion. The pH of the coarse emulsion was adjusted using sodium hydroxide. Following pH adjustment, the coarse emulsion was homogenized under high pressure to produce a fine particle size and thus a stable emulsion. The emulsion was filled into appropriate containers and sterilized in an autoclave.

[0058] These formulations were evaluated for microbial growth inhibition and the results are shown in Table 6 below.

Table 6

Formulation	Microbial Retardant (% w/v)	Time Point	EC	SA	PA	CA	AN
Example 2	0.005% EDTA	0	25	37	27	24	37
		24 hours	1	30	2	75	44
		48 hours	1	22	1	99	50
Example 8	0.025% EDTA	0	29	42	23	26	35
		24 hours	1	38	1	82	32
		48 hours	1	26	2	112	30
Example 9	0.001% EDTA	0	29	41	15	29	40
		24 hours	2	42	0	115	39
		48 hours	1	26	0	133	39
Example 10	0.0025% EDTA / 0.02% Sodium Citrate	0	30	45	7	29	38
		24 hours	2	32	2	68	37
		48 hours	1	29	0	85	88
Example 11	0.001% EDTA / 0.02% Sodium Citrate	0	26	32	29	22	40
		24 hours	1	39	0	64	41
		48 hours	1	26	7	40	41

[0059] The results demonstrate that varying concentrations of EDTA either alone or in combination with sodium citrate were effective in retarding microbial growth when added as a part of the manufacturing process.

[0060] Example 5 - The physical stability of emulsions with anti microbial agents was evaluated via zeta potential measurement. The zeta potential, which is a measure of charge around the oil droplet, was determined to evaluate the physical stability of emulsion formulations containing antimicrobial agents. A larger absolute value of zeta potential results in increased electrostatic repulsion between oil droplets and is indicative of enhanced physical stability. The results demonstrated that the absolute value of zeta potential decreased in presence of EDTA

alone or in combination with sodium citrate which was surprising given that their concentrations were low.

[0061] In an attempt to improve the physical stability of emulsions, use of oleic acid a co-emulsifier was investigated. Clevidipine emulsion with oleic acid as shown in the example below was manufactured using the following process: The oil phase containing soybean oil, clevidipine, egg yolk phospholipids and oleic acid was mixed with the aqueous phase containing glycerin, disodium edetate and water at approximately 70° C to form a coarse emulsion. The pH of the coarse emulsion was adjusted using sodium hydroxide. Following pH adjustment, the coarse emulsion was homogenized under high pressure to produce a fine particle size and thus a stable emulsion. The emulsion was filled into appropriate containers and sterilized in an autoclave.

Table 7

Function		% w/v
		Example 12
Active ingredient	Clevidipine butyrate	0.05
Oil phase	Soybean oil	20
Isotonicity adjuster	Glycerin	2.25
Microbial Growth Retardant	Disodium Edetate	0.005
Coemulsifier	Oleic acid	0.03
Emulsifier	Purified egg yolk phospholipids	1.2
pH adjuster	Sodium hydroxide	qs
Aqueous Phase	Water for Injection	qs to 100%

[0062] The above composition was evaluated for zeta potential and microbial growth inhibition.

[0063] Inhibition of microbial growth for Example 12 is shown in Table 8 below for 12, 24, and 30 hours.

Table 8

Challenge Organism	CFU/5mL			
	0 Hours	12 Hours	24 Hours	30 Hours
<i>S. marsacens</i> (SM)	29	32	32	40
<i>Staph. epidermidis</i> (SE)	30	6	5	5

[0064] The results for zeta potential are shown in Table 9 below and demonstrate that inclusion of oleic acid at a low concentration unexpectedly showed a substantial increase in zeta potential and physical stability of the emulsion compared to emulsions comprising EDTA alone or in combination with sodium citrate.

Table 9

Formulation	Microbial Retardant (% w/v)	Co-emulsifier (%w/v)	Zeta Potential (mV)	pH
Example 1	None	None	-25.6	7.7
Example 2	0.005% EDTA (Lab scale)	None	-24.8	7.7
Example 2	0.005% EDTA (pilot scale)	None	-17.6	7.7
Example 11	0.001% EDTA/ 0.02% SC	None	-18.3	7.4
Example 12	0.005% EDTA	0.03% Oleic acid	-36.1	8.0

[0065] In addition as demonstrated in Table 10 below, oleic acid did not impact the antimicrobial effectiveness of EDTA.

Table 10

Formulation	Microbial Retardant (% w/v)	Time Point	EC	SA	PA	CA	AN
Example 2	0.005% EDTA	0	25	37	27	24	37
		24 hours	1	30	2	75	44
		48 hours	1	22	1	99	50
Example 11	0.005% EDTA/ 0.03% Oleic Acid	0	39	32	35	70	50
		24 hours	1	49	49	305	75
		48 hours	0	41	70	395	40

Claims

1. A pharmaceutical formulation comprising,
 - (a) clevipidine, or a pharmaceutically acceptable salt or ester thereof, present at about 0.01 to about 1.0% w/v,
 - (b) an antimicrobial agent, present at about 0.001 to about 1.0% w/v,
 - (c) a lipid, present at about 2 to about 30% w/v,
 - (d) an emulsifier, present at about 0.2 to about 2.0% w/v,
 - (e) a tonicity modifier, present at about 2 to about 3% w/v, and
 - (f) water to 100%.
2. The pharmaceutical formulation of claim 1, wherein the antimicrobial agent is selected from the group consisting of benzyl alcohol, EDTA, sodium ascorbate, citric acid, and mixtures, derivatives, and salts thereof.
3. The pharmaceutical formulation of claim 1, wherein the lipid is selected from the group consisting of soybean oil, safflower seed oil, olive oil, cottonseed oil, sunflower oil, sesame oil, peanut oil, corn oil, medium chain triglycerides, triacetin, propylene glycol diesters, monoglycerides, and a mixture of two or more thereof.
4. The pharmaceutical formulation of claim 1, wherein the emulsifier is selected from the group consisting of egg yolk phospholipids, soybean phospholipids, synthetic phosphatidyl cholines, purified phosphatidyl cholines and hydrogenated phosphatidyl choline, and mixtures of two or more thereof.
5. The pharmaceutical formulation of claim 1, wherein the formulation has a pH of about 6.0 to about 8.8.
6. The pharmaceutical formulation of claim 1, wherein the formulation is contained in a sterile pre-filled syringe.
7. A pharmaceutical formulation comprising,
 - (a) clevipidine, or a pharmaceutically acceptable salt or ester thereof, present at about 0.01% to about 0.1% w/v,
 - (b) an antimicrobial agent, present at about 0.001 to about 1.0% w/v,
 - (c) a lipid, present at about 2 to about 30% w/v,
 - (d) an emulsifier, present at about 0.2 to about 2% w/v,

- (e) an antioxidant, present at about 0.01 to about 1.0% w/v,
 - (f) a tonicity modifier, present at about 2 to about 3% w/v, and
 - (g) water, to 100%.
8. The pharmaceutical formulation of claim 7, wherein the antimicrobial agent is selected from the group consisting of benzyl alcohol, EDTA, sodium ascorbate, citric acid, and mixtures, derivatives, and salts thereof.
9. The pharmaceutical formulation of claim 7, wherein the lipid is selected from the group consisting of soy bean oil, safflower seed oil, olive oil, cottonseed oil, sunflower oil, sesame oil, peanut oil, corn oil, medium chain triglycerides, triacetin, propylene glycol diesters, monoglycerides, and a mixture of two or more thereof.
10. The pharmaceutical formulation of claim 7, wherein the emulsifier is selected from the group consisting of egg yolk phospholipids, soybean phospholipids, synthetic phosphatidyl cholines, purified phosphatidyl cholines and hydrogenated phosphatidyl choline, and mixtures of two or more thereof.
11. The pharmaceutical formulation of claim 7, wherein the antioxidant is selected from the group consisting of sodium ascorbate, sodium citrate, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), propyl gallate, tocopherol, and a pharmaceutically acceptable salt thereof.
12. The pharmaceutical formulation of claim 7, wherein the formulation has a pH of about 6.0 to about 8.8.
13. The pharmaceutical formulation of claim 7, wherein the formulation is contained in a sterile pre-filled syringe.
14. A pharmaceutical formulation comprising,
- (a) clevidipine, or a pharmaceutically acceptable salt or ester thereof, present at about 0.01 to about 1% w/v,
 - (b) an antimicrobial agent, present at about 0.001 to about 1.0% w/v,
 - (c) a lipid, present at about 2 to about 30% w/v,
 - (d) an emulsifier, present at about 0.2 to about 2% w/v,
 - (e) an antioxidant, present at about 0.01 to about 1.0% w/v,
 - (f) a co-emulsifier, present at about 0.01 to about 2% w/v,

- (g) a tonicity modifier, present at about 2 to about 3% w/v, and
- (h) water, to 100%.

15. The pharmaceutical formulation of claim 14, wherein the antimicrobial agent is selected from the group consisting of benzyl alcohol, EDTA, sodium ascorbate, sodium citrate, and mixtures, derivates, and salts thereof.

16. The pharmaceutical formulation of claim 14, wherein the lipid is selected from the group consisting of soy bean oil, safflower seed oil, olive oil, cottonseed oil, sunflower oil, sesame oil, peanut oil, corn oil, medium chain triglycerides, triacetin, propylene glycol diesters, monoglycerides and a mixture of two or more thereof.

17. The pharmaceutical formulation of claim 14, wherein the emulsifier is selected from the group consisting of egg yolk phospholipids, soy bean phospholipids, synthetic phosphatidyl cholines, purified phosphatidyl cholines, hydrogenated phosphatidyl choline, and mixtures of two or more thereof.

18. The pharmaceutical formulation of claim 14, wherein the antioxidant is selected from the group consisting of sodium ascorbate, sodium citrate, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, tocopherol, and a pharmaceutically acceptable salt thereof.

19. The pharmaceutical formulation of claim 14, wherein the co-emulsifier is selected from the group consisting of glycerol, glycerin, poloxamers, Cremophor™, poloxamines, polyoxyethylene stearates, polyoxyethylene sorbitan fatty acid esters, sorbitan fatty acid esters, polysorbates, tocopherol PEG succinate, cholic acid, deoxycholic acid, oleic acid, and pharmaceutically acceptable salts thereof.

20. The pharmaceutical formulation of claim 14, wherein H324/78 is present in the formulation at about 0.2% or less.

21. The pharmaceutical formulation of claim 14, wherein the formulation has a pH of about 6.0 to about 8.8.


22. The pharmaceutical formulation of claim 14, wherein the formulation is contained in a sterile pre-filled syringe.

- 23. A pharmaceutical formulation comprising,
 - (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,

- (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) soybean oil, present at about 4 to about 30% w/v,
 - (d) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (e) glycerin, present at about 2 to about 3% w/v, and
 - (f) water up to 100%.
24. The pharmaceutical formulation of claim 23, wherein the formulation has a pH of about 6 to about 8.8.
25. A pharmaceutical formulation comprising,
- (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,
 - (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) soybean oil, present at about 4 to about 30% w/v,
 - (d) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (e) sodium citrate, present at about 0.005 to about 0.5% w/v,
 - (f) glycerin, present at about 2 to about 3% w/v, and
 - (g) water to 100% w/v.
26. The pharmaceutical formulation of claim 25, wherein the formulation has a pH of about 6 to about 8.8.
27. A pharmaceutical formulation comprising,
- (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,
 - (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) soybean oil, present at about 4 to about 30% w/v,
 - (d) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (e) sodium ascorbate, present at about 0.01 to about 1% w/v,
 - (f) glycerin, present at about 2 to about 3% w/v, and
 - (g) water, to 100%% w/v.
28. The pharmaceutical formulation of claim 27, wherein the formulation has a pH of about 6 to about 8.8.
29. A pharmaceutical formulation comprising,
- (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,
 - (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) oleic acid, present at about 0.01 to about 2.0% w/v,

- (d) soybean oil, present at about 4 to about 30% w/v,
 - (e) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (f) glycerin, present at about 2 to about 3% w/v, and
 - (g) water, up to 100% w/v.
30. The pharmaceutical formulation of claim 29, wherein the formulation has a pH of about 6 to about 8.8.
31. A pharmaceutical formulation comprising,
- (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,
 - (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) oleic acid, present at about 0.01 to about 2.0% w/v,
 - (d) soybean oil, present at about 4 to about 30% w/v,
 - (e) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (f) sodium citrate, present at about 0.005 to about 0.5% w/v,
 - (g) glycerin, present at about 2 to about 3% w/v, and
 - (h) water, up to 100% w/v.
32. The pharmaceutical formulation of claim 31, wherein the formulation has a pH of about 6 to about 8.8.
33. A pharmaceutical formulation comprising,
- (a) clevidipine butyrate, present at about 0.01 to about 1% w/v,
 - (b) EDTA, present at about 0.001 to about 0.1% w/v,
 - (c) oleic acid, present at about 0.01 to about 2% w/v,
 - (d) soybean oil, present at about 4 to about 30% w/v,
 - (e) purified egg yolk phospholipids, present at about 0.2 to about 2% w/v,
 - (f) sodium ascorbate, present at about 0.05 to about 1.0% w/v,
 - (g) glycerin, present at about 2 to about 3% w/v, and
 - (h) water, up to 100%.
34. The pharmaceutical formulation of claim 33, wherein the formulation has a pH of about 6 to about 8.5.

Figure 1. Clevidipine Degradation Pathway



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 11/55617

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - A01N 37/30; A61K 31/205 (2012.01)
 USPC - 514/356

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 USPC - 514/356

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 USPC - 424/400, 424/489; 546/321; 560/51; 562/459 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 *** Databases: WEST (PGPB, USPT, USOC, EPAB, JPAB); Google, Google Scholar *** Search Terms Used: Medicines Company, Motheram, Williams, clevidipine, cleviprex, dihydropyridine calcium channel blocker, antimicrobial, EDTA, benzyl alcohol, sodium ascorb

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2010/0113534 A1 (Motheram et al.) 06 May 2010 (06.05.2010) especially para [0002]-[0003], [0008], [0011], [0028], [0041], [0051]-[0053]	1-34
Y	US 2006/0104913 A1 (Chaudry) 18 May 2006 (18.05.2006) especially para [0036], [0038], [0044]-[0049], [0052], [0089]	1-34
A	WO 2010/022259 A1 (Maguire et al.) 25 February 2010 (25.02.2010) entire document	1-34
A	US 2010/0105743 A1 (Krishna et al.) 29 April 2010 (29.04.2010) entire document	1-34

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
 14 February 2012 (14.02.2012)

Date of mailing of the international search report
08 MAR 2012

Name and mailing address of the ISA/US
 Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, Virginia 22313-1450
 Facsimile No. 571-273-3201

Authorized officer:
 Lee W. Young
 PCT Helpdesk: 571-272-4300
 PCT OSP: 571-272-7774