

Mini- and microsatellite expansions: the recombination connection

Guy-Franck Richard^{1,+} and Frédéric Pâques^{2,+}

¹Unité de Génétique Moléculaire des Levures, URA 2171 CNRS and UFR 927 Université Pierre et Marie Curie, Paris, Institut Pasteur, 25 rue du Dr Roux, 75724 Paris Cedex 15 and ²Institut Jacques Monod, UMR 7592, Université Paris 6 et Paris 7, Tour 43, 2 Place Jussieu, 75251 Paris Cedex 05, France

Received May 24, 2000; revised June 21, 2000; accepted June 28, 2000

It is widely accepted that the large trinucleotide repeat expansions observed in many neurological diseases occur during replication. However, genetic recombination has emerged as a major source of instability for tandem repeats, including minisatellites, and recent studies raise the possibility that it may also be responsible for trinucleotide repeat expansions. We will review data connecting tandem repeat rearrangements and recombination in humans and in eukaryotic model organisms, and discuss the possible role of recombination in trinucleotide repeat expansions in human neurological disorders.

Introduction

Two classes of tandem repeat sequences, minisatellites and microsatellites, have gained increasing attention from the scientific community over the past decade. Microsatellites are tandem arrays of short (usually <10 bp) units, while minisatellites are tandem arrays of longer units (>10 and <100 bp). These two kinds of sequence are widespread in all eukaryotic genomes (reviewed in Charlesworth *et al.*, 1994) from yeast to mammals. A growing number of neurological disorders have been found to result from the expansion of a particular class of microsatellites, called trinucleotide repeats (reviewed in Cummings and Zoghbi, 2000).

In contrast to other forms of microsatellite instability such as that found in mismatch repair (MMR) mutants, trinucleotide repeat expansions observed in neurological diseases are often quite large (several times the initial number of repeats). The occurrence of these expansions during the lifetime is not precisely known; they might be meiotic, pre- or post-zygotic. Trinucleotide repeats are not the only class of tandem repeats prone to expansions. Three minisatellites are known to share this property. Expansion of one of them is associated with a human

disease (Lalioi *et al.*, 1997). The others, two AT-rich repeats, are responsible for fragile sites (Yu *et al.*, 1997; Hewett *et al.*, 1998).

Two types of mechanism have been proposed to be involved in tandem repeat instability: replication and recombination. The genome-wide microsatellite instability observed in MMR mutant cells (Simpkins *et al.*, 1999 and references therein) strongly supports a replicative mechanism, and the somatic instability of trinucleotide repeats is generally viewed as a consequence of replication (reviewed in McMurray, 1999; Richard *et al.*, 1999a). However, several recent results argue that recombination might also play a role in microsatellite expansions. The aim of this paper is to review the data connecting tandem repeat rearrangements with recombination, and to discuss the possible involvement of recombination in trinucleotide repeat expansions.

Gene conversion as a major source of tandem repeat instability

Initially, when people were thinking of tandem repeat rearrangements in terms of recombination, they were thinking of crossovers (Figure 1), i.e. a reciprocal transfer of genetic information. However, gene conversion (non-reciprocal transfer of information) has recently emerged as a major cause of tandem repeat instability. Gene conversion and crossover are often but not always associated, and a series of models, culminating in the model proposed by Szostak *et al.* (1983) (see Figure 2A), summarizes this relationship. However, it is mostly gene conversion without crossover that destabilizes tandem repeats.

Welch *et al.* (1990) were the first to establish a correlation between non-crossover gene conversion and tandem repeat rearrangements, when they observed meiotic expansions and contractions of the yeast *CUP1* locus. Tandem repeat rearrangements that were likely to have resulted from gene conversion were also

*Corresponding authors. G.-F.R.: Tel: +33 1 40 61 34 54; Fax: +33 1 40 61 34 56; E-mail: gfrichar@pasteur.fr
F.P.: Tel: +33 1 44 27 57 07; Fax: +33 1 44 27 52 65; E-mail: paques@ijm.jussieu.fr




Fig. 1. Tandem repeat rearrangement by unequal crossover between sister chromatids. Reciprocal exchange leads to simultaneous contraction of one of the repeats and expansion of the other.

observed in *Drosophila* (Pâques and Wegnez, 1993; Thompson-Stewart *et al.*, 1994; Delattre *et al.*, 1995; Pâques *et al.*, 1996) and provided the basis for a new model for tandem repeat rearrangements. These events were thought to result from the excision of a *Drosophila* P transposon. Such excisions are repaired by gene conversion, which is not usually associated with crossovers. A second series of models, termed SDSA, for synthesis-dependent strand annealing (reviewed in Pâques and Haber, 1999), accounts for non-crossover gene conversions. In SDSA models, the two newly synthesized strands unwind from their template and anneal with each other (Figure 2B). Pâques and Wegnez (1993) postulated that out-of-frame re-annealing of tandem repeats would explain the observed rearrangements detected after excision of a P transposon (Figure 2C).

Meiotic recombination in human minisatellites

Since the discovery of the first minisatellites in the human genome (Jeffreys *et al.*, 1985), mechanisms leading to high spontaneous mutation rates of these hypervariable DNA sequences have been thoroughly studied. Jeffreys *et al.* (1988) proposed that small mutations involving gain or loss of 4–10 repeats could occur by a mitotic replication slippage mechanism, while large mutation events involving gain or loss of up to 200 repeats were more compatible with a meiotic recombinational process.

In further analyses, minisatellite mutations found in the germline were precisely mapped (Buard and Vergnaud, 1994; Jeffreys *et al.*, 1994), and typical features of the mutation process were observed. First, there was a polarity in the mutational process, i.e. repeats were gained or lost preferentially at one end of the repeat array. Secondly, complex rearrangement events suggested that the new mutant allele gained genetic information from both the sister chromatid (intra-allelic event) and the homologous chromosome (inter-allelic event). Thirdly, exchange of flanking markers was rare, indicating that recombination occurred with few associated crossovers.

Again, these results were explained in terms of SDSA (Buard and Jeffreys, 1997). This mechanism provided a simple explanation for the complex events observed: both templates were used by the two ends of a double-strand break (DSB) to copy information, and the two newly synthesized strands were assembled in the recipient molecule by annealing. SDSA could also

explain the low crossover rate (Figure 2B). Related SDSA models, where DNA synthesis occurred at only one end of the DSB, may also account for other features of minisatellite rearrangements observed in yeast or humans (Pâques *et al.*, 1998; Debrauwère *et al.*, 1999).

By mapping these rare crossovers in the vicinity of the MS32 minisatellite, Jeffreys *et al.* (1998) found a meiotic hot spot (a chromosomal region where meiotic recombination is frequently observed) centered ~200 bp upstream of MS32 and extending into the beginning of the repeat array. Their work strongly suggests that hypervariability arises in proximity to a meiotic hot spot. Their finding also explains mutational polarity, since meiotic DSBs are initiated on one side of the minisatellite and recombination propagates within the repeat array from the initiation point. The gene conversion gradient decreases from the initiation point, as observed for meiotic hot spots in yeast (Nicolas *et al.*, 1989).

Analysis of somatic mutations at MS32 revealed simple deletions or duplications, with no detectable bias toward one extremity of the repeat array (Jeffreys and Neumann, 1997). The authors thus concluded that somatic mutational processes involved in MS32 expansions and contractions were different from those found in the germline.

The introduction of two minisatellites, MS32 and CEB1, into yeast confirmed that they are more unstable during meiosis than during mitosis. Gains and losses of repeats were again complex events involving both inter- and intra-allelic recombination (Appelgren *et al.*, 1997; Debrauwère *et al.*, 1999). In addition, it was shown that CEB1 meiotic instability depended on DSB formation at two well characterized meiotic hot spots, one of which lies upstream and the other downstream of the repeat array. Interestingly, instability disappeared when CEB1 was moved to a chromosomal region where no meiotic DSB could be detected.

Gene conversion-induced rearrangements of tandem repeats in yeast

Recently, Pâques *et al.* (1998) designed an experimental system to study the effect of mitotic gene conversion on tandem repeat instability in *Saccharomyces cerevisiae*. A DSB was made on a yeast chromosome, which could be repaired using a homologous sequence containing a tandem repeat as donor (Figure 3). Both




Fig. 2. Recombination models. In all models, gene conversion is initiated by a DSB, and strand invasion of the homologous template is initiated by a resected 3' end. Arrows represent DNA synthesis associated with DSB-repair. The broken molecule ('donor') is shown in blue, the template molecule ('recipient') is shown in red, newly synthesized DNA is shown in orange. Tandem repeats are represented by hatched lines. (A) Szostak *et al.* (1983) model. Resolution of the two four-stranded structures called 'Holliday junctions' (circled in black) results in gene conversion associated with crossovers in 50% of the cases. (B) SDSA model. Resolution is achieved by unwinding the two newly synthesized strands from the template and annealing them. No crossover is expected. (C) SDSA model of tandem repeat rearrangements. The newly synthesized strands both contain repeats and their out-of-frame annealing can lead to contraction or expansion of the tandem repeat. (C') Alternatively, contraction and expansion can result from unwinding and out-of-frame re-invasion of one (or both) the newly synthesized strand(s). Only one kind of event (expansion) is shown here. Several rounds of unwinding/re-invasion could take place during a single gene conversion, leading to successive rounds of DNA synthesis.

donor and recipient were recovered and analyzed for tandem repeat rearrangements.

Different kinds of tandem repeat were studied including 5S rRNA genes, trinucleotide repeats and minisatellites (Pâques *et al.*, 1998; Richard *et al.*, 1999b, 2000; F. Pâques, G.-F. Richard and J.E. Haber, unpublished results), all of which rearranged with a high frequency during gene conversion (Table I). Furthermore, rearrangements were observed mainly in the recipient

molecule, whereas the donor was extremely stable, as predicted by SDSA models (Figure 2C). Interestingly, the CAG₃₉ repeat (i.e. an array of 39 CAG triplets) yielded only contractions, whereas both contractions and expansions were observed with CAG₉₈ (Table I).

These results demonstrate the destabilizing effect of gene conversion on different kinds of tandem repeat. The model proposed in Figure 2C postulates that, during gene conversion-associated DNA synthesis, slippage of the newly synthesized strands can occur, leading to addition or subtraction of a whole number of repeats. The frequency of such 'repair-slippage' was estimated to be 800-fold higher than that for S-phase replication-slippage (Richard *et al.*, 1999b).

Is DSB-repair involved in trinucleotide repeat expansions?

Trinucleotide repeats and some minisatellites are fragile sites in human (reviewed in Sutherland *et al.*, 1998) and in yeast (Freudenreich *et al.*, 1998). Fragile sites are loci at which chromosome breakage occurs in the presence of specific chemicals. We suggest that a mitotic or meiotic DSB could initiate the expansion process. Meiotic instability of CAG repeats in yeast has been correlated with formation of a nearby DSB in one study using long CAG tracts (Jankowski *et al.*, 2000), but not in another using shorter CAG tracts (Moore *et al.*, 1999). White *et al.* (1999) showed that CCG repeats did not exhibit a higher rate of instability during meiosis than during vegetative growth in yeast, whereas Cohen *et al.* (1999) found that instability of CTG repeats carried by YACs was higher during meiosis than during mitotic divisions. A very recent study showed that CA dinucleotide repeats were also destabilized during meiosis, in yeast, and that progression of strand exchange during gene conversion was inhibited by the minisatellite (Gendrel *et al.*, 2000). In transgenic mice, CAG repeats exhibited a higher instability in the germline than in somatic cells, with a propensity towards expansion rather than contraction (Seznec *et al.*, 2000).

In human diseases, it is common to observe expansions of more than twice the original size. Such large expansions in yeast and humans could occur by successive rounds of unwinding/re-invasion of the donor sequence by the newly synthesized strand, allowing DNA synthesis to proceed more than once within the repeats (Pâques *et al.*, 1998; Figure 2C').

An interesting case of large contraction of a CAG repeat was described during transmission of a myotonic dystrophy allele. O'Hoy *et al.* (1993) reported a large reduction of the number of CAG triplets associated with what they called a 'discontinuous gene conversion event'. The resulting allele was a patchwork of both maternal and paternal alleles. Buard and Vergnaud (1994), Debrauwère *et al.* (1999) and Tamaki *et al.* (1999) also found complex recombination events in minisatellites. In those cases, gene conversion was inter-allelic (between homologous chromosomes), intra-allelic (between sister chromatids) or a patchwork of both. All these observations are compatible with the unwinding/re-invasion model (Figure 2C').

Perspectives

Even though more conclusive evidence will be required to account for dramatic trinucleotide repeat expansion by a gene




Fig. 3. Experimental system used to study tandem repeat instability during mitotic gene conversion induced by an endonuclease in yeast. After induction of a DSB on a yeast chromosome by an inducible endonuclease, yeast cells can repair the broken molecule using a tandem repeat-containing homologous donor sequence. During DNA synthesis associated with DSB-repair, slippage of the newly synthesized strand(s) may occur, giving rise to contractions or expansions of the tandem repeat in the recipient molecule. The blue circle represents the chromosome centromere.

Table I. Rearrangements of tandem repeat sequences during gene conversion

	Faithful gene conversion (%)	Gene conversion with contraction (%)	Gene conversion with expansion (%)	Total (contractions + expansions) (%)
5S rDNA genes	63	23	14	37
CAG ₃₉	84	16	–	16
CAG ₉₈	57	30	13	43
CAA ₈₇	90	10	–	10

conversion mechanism in humans, recent data suggest that recombination can lead to large scale contractions and expansions within tandem repeat sequences at a high frequency. Experimental systems can be designed to study the effect of mitotic and meiotic recombination on tandem repeat instability. It will be particularly interesting to determine whether any kind of sequence can be expanded during gene conversion, or whether there is a sequence (or structure) specificity to this process. Moreover, even though some *trans*-acting factors have been shown to be involved in expansions during gene conversion (Richard *et al.*, 2000), their precise function needs to be determined. Hopefully, further experiments in model organisms will allow us to unravel the molecular basis of tandem repeat complex rearrangements during gene conversion, and perhaps help us to understand how dramatic trinucleotide repeat expansions occur in humans.

Acknowledgements

We would like to thank Allyson Holmes and members of the Unité de Génétique Moléculaire des Levures for insightful suggestions. G.-F.R. is the recipient of a post-doctoral fellowship from the Institut Pasteur.

References

- Appelgren, H., Cederberg, H. and Rannug, U. (1997) Mutations at the human minisatellite MS32 integrated in yeast occur with high frequency in meiosis and involve complex recombination events. *Mol. Gen. Genet.*, **256**, 7–17.
- Buard, J. and Jeffreys, A.J. (1997) Big, bad minisatellites. *Nature Genet.*, **15**, 327–328.
- Buard, J. and Vergnaud, G. (1994) Complex recombination events at the hypermutable minisatellite CEB1 (D2S90). *EMBO J.*, **13**, 3203–3210.
- Charlesworth, B., Sniegowski, P. and Stephan, W. (1994) The evolutionary dynamics of repetitive DNA in eukaryotes. *Nature*, **371**, 215–220.
- Cohen, H., Sears, D.D., Zenvirth, D., Hieter, P. and Simchen, G. (1999) Increased instability of human CTG repeat tracts on yeast artificial chromosomes during gametogenesis. *Mol. Cell. Biol.*, **19**, 4153–4158.
- Cummings, C.J. and Zoghbi, H.Y. (2000) Fourteen and counting: unraveling trinucleotide repeat diseases. *Hum. Mol. Genet.*, **9**, 909–916.
- Debrauwère, H., Buard, J., Tessier, J., Aubert, D., Vergnaud, G. and Nicolas, A. (1999) Meiotic instability of human minisatellite CEB1 in yeast requires double-strand breaks. *Nature Genet.*, **23**, 367–371.
- Delattre, M., Anxolabehere, D. and Coen, D. (1995) Prevalence of localized rearrangements vs. transpositions among events induced by *Drosophila* P element transposase on a P transgene. *Genetics*, **141**, 1407–1424.
- Freudenreich, C.H., Kantrow, S.M. and Zakian, V.A. (1998) Expansion and length-dependent fragility of CTG repeats in yeast. *Science*, **279**, 853–856.

G.-F. Richard and F. Pâques

- Gendrel, C.-G., Boulet, A. and Dutreix, M. (2000) (CA/GT)_n microsatellites affect homologous recombination during yeast meiosis. *Genes Dev.*, **14**, 1261–1268.
- Hewett, D.R. *et al.* (1998) *FRA10B* structure reveals common elements in repeat expansion and chromosomal fragile site genesis. *Mol. Cell*, **1**, 773–781.
- Jankowski, C., Nasar, F. and Nag, D.K. (2000) Meiotic instability of CAG repeat tracts occurs by double-strand break repair in yeast. *Proc. Natl Acad. Sci. USA*, **97**, 2134–2139.
- Jeffreys, A.J. and Neumann, R. (1997) Somatic mutation processes at a human minisatellite. *Hum. Mol. Genet.*, **6**, 129–136.
- Jeffreys, A.J., Wilson, V. and Thein, S.L. (1985) Hypervariable ‘minisatellite’ regions in human DNA. *Nature*, **314**, 67–73.
- Jeffreys, A.J., Royle, N.J., Wilson, V. and Wong, Z. (1988) Spontaneous mutation rates to new length alleles at tandem-repetitive hypervariable loci in human DNA. *Nature*, **332**, 278–281.
- Jeffreys, A.J., Tamaki, K., McLeod, A., Monckton, D.G., Neil, D.L. and Armour, J.A.L. (1994) Complex gene conversion events in germline mutation at human minisatellites. *Nature Genet.*, **6**, 136–145.
- Jeffreys, A.J., Murray, J. and Neumann, R. (1998) High-resolution mapping of crossovers in human sperm defines a minisatellite-associated recombination hot spot. *Mol. Cell*, **2**, 267–273.
- Lalioti, M.D., Scott, H.S., Buresi, C., Rossier, C., Bottani, A., Morris, M.A., Malafose, A. and Antonarakis, S.E. (1997) Dodecamer repeat expansion in cystatin B gene in progressive myoclonus epilepsy. *Nature*, **386**, 847–851.
- McMurray, C.T. (1999) DNA secondary structure: a common and causative factor for expansion in human disease. *Proc. Natl Acad. Sci. USA*, **96**, 1823–1825.
- Moore, H., Greenwell, P.W., Liu, C.-P., Arnheim, N. and Petes, T.D. (1999) Triplet repeats form secondary structures that escape DNA repair in yeast. *Proc. Natl Acad. Sci. USA*, **96**, 1504–1509.
- Nicolas, A., Treco, D., Schultes, N.P. and Szostak, J.W. (1989) An initiation site for meiotic gene conversion in the yeast *Saccharomyces cerevisiae*. *Nature*, **338**, 35–39.
- O’Hoy, K.L., Tsilfidis, C., Mahadevan, M.S., Neville, C.E., Barcelo, J., Hunter, A.G. and Korneluk, R.G. (1993) Reduction in size of the myotonic dystrophy trinucleotide repeat mutation during transmission. *Science*, **259**, 809–812.
- Pâques, F. and Haber, J.E. (1999) Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiol. Mol. Biol. Rev.*, **63**, 349–404.
- Pâques, F. and Wegnez, M. (1993) Deletions and amplifications of tandemly arranged ribosomal 5S genes internal to a P element occur at a high rate in a dysgenic context. *Genetics*, **135**, 469–476.
- Pâques, F., Bucheton, B. and Wegnez, M. (1996) Rearrangements involving repeated sequences within a P element preferentially occur between units close to the transposon extremities. *Genetics*, **142**, 459–470.
- Pâques, F., Leung, W.-Y. and Haber, J.E. (1998) Expansions and contractions in a tandem repeat induced by double-strand break repair. *Mol. Cell. Biol.*, **18**, 2045–2054.
- Richard, G.-F., Hennequin, C., Thierry, A. and Dujon, B. (1999a) Trinucleotide repeats and other microsatellites in yeasts. *Res. Microbiol.*, **150**, 589–602.
- Richard, G.-F., Dujon, B. and Haber, J. (1999b) High frequency of rearrangements of short CAG/CTG trinucleotide repeats in yeast induced by double-strand break repair. *Mol. Gen. Genet.*, **261**, 871–882.
- Richard, G.-F., Goellner, G.M., McMurray, C.T. and Haber, J.E. (2000) Recombination-induced CAG trinucleotide repeat expansions in yeast involve the MRE11/RAD50/XRS2 complex. *EMBO J.*, **19**, 2381–2390.
- Seznec, H., Lia-Baldini, A.-S., Duros, C., Fouquet, C., Lacroix, C., Hofmann-Radvanyi, H., Junien, C. and Gourdon, G. (2000) Transgenic mice carrying large human genomic sequences with expanded CTG repeat mimic closely the DM CTG repeat intergenerational and somatic instability. *Hum. Mol. Genet.*, **9**, 1185–1194.
- Simpkins, S.B., Bocker, T., Swisher, E.M., Mutch, D.G., Gersell, D.J., Kovatich, J.P., Fishel, R. and Goodfellow, P.J. (1999) MLH1 promoter methylation and gene silencing is the primary cause of microsatellite instability in sporadic endometrial cancers. *Hum. Mol. Genet.*, **8**, 661–666.
- Sutherland, G.R., Baker, E. and Richards, R.I. (1998) Fragile sites still breaking. *Trends Genet.*, **14**, 501–506.
- Szostak, J.W., Orr-Weaver, T.L., Rothstein, R.J. and Stahl, F.W. (1983) The double-strand break repair model for recombination. *Cell*, **33**, 25–35.
- Tamaki, K., May, C.A., Dubrova, Y.E. and Jeffreys, A.J. (1999) Extremely complex repeat shuffling during germline mutation at human minisatellite B6.7. *Hum. Mol. Genet.*, **8**, 879–888.
- Thompson-Stewart, D., Karpen, G.H. and Spradling, A.C. (1994) A transposable element can drive the concerted evolution of tandemly repetitive DNA. *Proc. Natl Acad. Sci. USA*, **91**, 9042–9046.
- Welch, J.W., Maloney, D.H. and Fogel, S. (1990) Unequal crossing-over and gene conversion at the amplified *CUP1* locus of yeast. *Mol. Gen. Genet.*, **222**, 304–310.
- White, P.J., Borts, R.H. and Hirst, M.C. (1999) Stability of the human Fragile X (CCG)_n triplet repeat array in *Saccharomyces cerevisiae* deficient in aspects of DNA metabolism. *Mol. Cell. Biol.*, **19**, 5675–5684.
- Yu, S. *et al.* (1997) Human chromosomal fragile site *FRA16B* is an amplified AT-rich minisatellite repeat. *Cell*, **88**, 367–374.

DOI: 10.1093/embo-reports/kvd031